

JOURNAL OF NUMBER THEORY **39**, 225–236 (1991)

A Linear Independence Measure for Certain p -Adic Numbers

K. VÄÄNÄNEN*

*Department of Mathematics,
University of Oulu, 90570 Oulu 57, Finland*

AND

R. WALLISSER

*Mathematisches Institut der Albert-Ludwigs-Universität,
Albertstraße 23, 7800 Freiburg im Breisgau, Germany*

Communicated by W. Schmidt

Received May 2, 1989; revised March 5, 1990

DEDICATED TO THE MEMORY OF THEODOR SCHNEIDER

A linear independence measure is obtained for certain values of a p -adic function $\sum_{n=0}^{\infty} q^{n(n-1)/2} z^n$, where $q \in \mathbb{Q}$, $0 < |q|_p < 1$.

© 1991 Academic Press, Inc.

1. INTRODUCTION

In 1921 Tschakaloff [10] proved the linear independence over \mathbb{Q} of certain values of the function

$$f(z) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} z^n, \quad (1)$$

where q , $0 < |q| < 1$, is a rational number satisfying certain conditions. In the case $q^{-1} \in \mathbb{Z}$ these results were generalized by Skolem [7] to give the linear independence of the values of the derivatives of f . He used a method following closely certain transcendence proofs of e , see, e.g., Perron [6]. More recently the arithmetic properties of the values of f have been studied

* This work was carried out while the first of the authors was a research fellow of the Alexander von Humboldt Foundation.

by several authors; in particular the recent work of Bezzin [2] contains as a special case the above results. Further, quantitative irrationality measures for certain sets of values of f have been obtained by Bundschuh and Shiokawa [3] and Stihl [8]. In [11] we showed that Skolem's method can be refined to obtain irrationality measures for sets containing values of the derivatives of f .

The first studies on arithmetic properties of the p -adic function (1) in the case $q = p^m$, $m \in \mathbb{N}$, were made by Icen [4], who proved that this function cannot have algebraic values at too many algebraic points. The linear independence of certain values of this same p -adic function and its derivatives was obtained by Bezzin [2]. In [11] it was shown that Skolem's method can be used to obtain quantitative estimates also in the p -adic case if q , $0 < |q|_p < 1$, satisfies certain conditions. However, it seems that, e.g., the case $q = p$ cannot be handled by this method. In the present paper our purpose is to obtain a linear independence measure for the values of the p -adic function (1) by using the Padé approximation polynomials of Stihl [8]. In this paper we give a p -adic analogue of the main result of Bundschuh and Shiokawa [3] and of certain results of Stihl [8].

2. RESULTS

We shall use $|\cdot|$ to denote the usual archimedean absolute value. If p is a prime, then $|\cdot|_p$ is used to denote the p -adic valuation of \mathbb{Q}_p normalized by $|p|_p = p^{-1}$. Let us suppose that $q = s/r \in \mathbb{Q}$, $(s, r) = 1$, $r > 0$, and $|s| > 1$. If p is a prime factor of s , then $|q|_p < 1$ and the series (1) converges for all $z \in \mathbb{Q}_p$. Our main result is the following linear independence measure for some values of this p -adic function f .

THEOREM 1. *Let $\gamma_1, \dots, \gamma_D$ denote non-zero rational numbers such that $\gamma_i/\gamma_j \neq q^n$ for all $i \neq j$, $n \in \mathbb{Z}$. Suppose that*

$$\begin{aligned} \gamma &= 1 + (\log |s|_p) / \log \max\{|s|, r\} < \Gamma = \Gamma(D) \\ &= (2D + 1 - (1 + 4D^2)^{1/2}) / (2D). \end{aligned} \quad (2)$$

If $\varepsilon > 0$ is given, then there exists a positive constant C , depending on q , p , D , $\gamma_1, \dots, \gamma_D$ and ε , such that for any rational integers H_0, H_1, \dots, H_D , not all zero, we have

$$\left| H_0 + \sum_{j=1}^D H_j f(\gamma_j) \right|_p > CH^{-\theta-\varepsilon},$$

where $H = \max\{|H_0|, |H_1|, \dots, |H_D|\}$, and

$$\theta = \theta(\gamma, D) = 1 + (2D - 1 + (1 + 4D^2)^{1/2}) / (2 - \gamma(2D + 1 + (1 + 4D^2)^{1/2})).$$

We note that in the case $q = p^m$, $m \in \mathbb{N}$, γ has the value zero and thus (2) holds. This leads immediately to the following corollary concerning the values of the p -adic function

$$g(z) = \sum_{n=0}^{\infty} p^{an^2 + bn + c} z^n, \quad a, b, c \in \mathbb{Z}, a > 0.$$

COROLLARY. *Let $\gamma_1, \dots, \gamma_D$ denote non-zero rational numbers such that $\gamma_i/\gamma_j \neq p^{2an}$ for all $i \neq j$, $n \in \mathbb{Z}$. If $\varepsilon, H_0, \dots, H_D$ are as in Theorem 1, then there exists a positive constant C_1 , depending on $p, D, \gamma_1, \dots, \gamma_D, a, b, c$, and ε , such that*

$$\left| H_0 + \sum_{j=1}^D H_j g(\gamma_j) \right|_p > C_1 H^{-\theta - \varepsilon},$$

where $\theta = \theta(D) = (2D + 1 + (1 + 4D^2)^{1/2})/2$.

If s has several prime factors p , then it may happen that (2) does not hold for any of these p . The case $D = 1$, $q = 60$ is such an example. In this situation we are only able to prove the following result on the non-existence of “ s -global” relations.

THEOREM 2. *Let $\gamma_1, \dots, \gamma_D$ be as in Theorem 1. Suppose that*

$$1 - (\log |s|)/\log \max\{|s|, r\} < \Gamma.$$

If H_0, H_1, \dots, H_D are rational integers, not all zero, then the equality

$$H_0 + H_1 f_p(\gamma_1) + \dots + H_D f_p(\gamma_D) = 0$$

cannot hold for all $p|s$. Here, for each $p|s$, $f_p(\gamma_i)$ denotes the value $f(\gamma_i)$ of f in the corresponding \mathbb{Q}_p .

3. PADÉ APPROXIMATION POLYNOMIALS

Recently Stihl [8] used Maier’s [5] method to find Padé approximation polynomials of the second kind for the functions $f(\gamma_1 z), \dots, f(\gamma_D z)$. The considerations of Stihl can be applied also in the p -adic case if $|q|_p < 1$. Thus Satz 4 of [8] implies the following lemma (we choose there $P(x) = x$, $Q(x) \equiv 1$).

LEMMA 1. *Let $\gamma_1, \dots, \gamma_D$ be rational numbers as in Theorem 1. For the parameters $l, \lambda \in \mathbb{N}$ we define the polynomial $A_{l,\lambda}$ by*

$$A_{l,\lambda}(z) = \sum_{i=0}^{Dl} z^{Dl-i} \sigma_{i,l} q^{-\binom{i+\lambda}{2}},$$

where $\sigma_{i,t}$ are defined by the identity

$$\prod_{u=1}^D \prod_{k=0}^{l-1} (\gamma_u - wq^{-k}) = \sum_{i=0}^{Dl} \sigma_{i,t} w^i.$$

Further, let

$$B_{l,\lambda,t}(z) = \sum_{k=0}^{Dl+\lambda-1} E_{k,t} z^k, \quad t = 1, \dots, D,$$

where the coefficients $E_{k,t}$ are given by

$$E_{k,t} = \sum_{\substack{i=0 \\ i+k-Dl \geq 0}}^{Dl} \sigma_{i,t} \gamma_t^{i+k-Dl} q^{\binom{i+k-Dl}{2} - \binom{i+\lambda}{2}}, \\ k = 0, 1, \dots, t = 1, \dots, D.$$

Then we have, for all $t = 1, \dots, D$,

$$A_{l,\lambda}(z) f(\gamma_t z) - B_{l,\lambda,t}(z) = \sum_{k=l+Dl+\lambda}^{\infty} E_{k,t} z^k = R_{l,\lambda,t}(z).$$

The proof of our results is based on certain important properties of the polynomials $A_{l,\lambda}$ and $B_{l,\lambda,t}$ which we shall consider in the following paragraph.

4. ON PROPERTIES OF $A_{l,\lambda}$ AND $B_{l,\lambda,t}$

In our considerations we shall use, for all $a, b, c \in \mathbb{Q}$, the notation

$$(a+b, c)_0 = 1, \quad (a+b, c)_l = \prod_{k=0}^{l-1} (a+bc^k), \quad l = 1, 2, \dots$$

From [1, p. 36], it follows that for all $l \in \mathbb{N}$

$$(a+b, c)_l = \sum_{k=0}^l \begin{bmatrix} l \\ k \end{bmatrix}_c c^{\binom{k}{2}} a^{l-k} b^k, \quad (3)$$

where, for all $k = 0, 1, \dots, l$,

$$\begin{bmatrix} l \\ k \end{bmatrix}_c = (1-c, c)_l / ((1-c, c)_k (1-c, c)_{l-k}).$$

We begin by proving a lemma which gives a common denominator for the coefficients of the polynomials $A_{l,\lambda}$ and $B_{l,\lambda,t'}$ and an upper estimate for

the absolute values at $z=1$ of these polynomials. By c_1, c_2, \dots we shall denote effectively computable positive constants independent of l and λ .

LEMMA 2. Let $h = h(q) = \max\{|s|, r\}$, and let us put $\gamma_i = a_i/b_i$, $(a_i, b_i) = 1$, $i = 1, \dots, D$. Further, let $\Omega_{l,\lambda}$ denote a natural number

$$\Omega_{l,\lambda} = s^{e(l,\lambda)} (b_1 \dots b_D)^{(D+1)l + \lambda - 1},$$

where $e(l, \lambda) = [(l^2(D^2 + D) + \lambda(2Dl + \lambda))/2]$. Then all of the numbers

$$\Omega_{l,\lambda} A_{l,\lambda}(1), \quad \Omega_{l,\lambda} B_{l,\lambda,t}(1), \quad t = 1, \dots, D,$$

are rational integers satisfying the inequality

$$\max\{|\Omega_{l,\lambda} A_{l,\lambda}(1)|, |\Omega_{l,\lambda} B_{l,\lambda,t}(1)|\} \leq c_1^{l+\lambda} h^{e(l,\lambda)}.$$

Proof. By (3) and the definition of $\sigma_{i,l}$ we obtain

$$\begin{aligned} \sum_{i=0}^{Dl} \sigma_{i,l} w^i &= \prod_{u=1}^D (\gamma_u - w, q^{-1})_l \\ &= \prod_{u=1}^D \left(\sum_{k=0}^l \begin{bmatrix} l \\ k \end{bmatrix}_{1/q} q^{-\binom{k}{2}} \gamma_u^{l-k} (-w)^k \right) \\ &= \sum_{i=0}^{Dl} (-1)^i \sum_{\substack{v_1 + \dots + v_D = i \\ 0 \leq v_j \leq l}} \\ &\quad \times \left\{ \begin{bmatrix} l \\ v_1 \end{bmatrix}_{1/q} \dots \begin{bmatrix} l \\ v_D \end{bmatrix}_{1/q} q^{-\binom{v_1}{2} - \dots - \binom{v_D}{2}} \gamma_1^{l-v_1} \dots \gamma_D^{l-v_D} w^i \right\}. \end{aligned}$$

Since

$$\begin{bmatrix} l \\ v \end{bmatrix}_{1/q} q^{-\binom{v}{2}} = q^{-vl} \begin{bmatrix} l \\ v \end{bmatrix}_q q^{\binom{v+1}{2}},$$

this yields

$$\begin{aligned} \sigma_{i,l} &= (-1)^i q^{-il} \sum_{\substack{v_1 + \dots + v_D = i \\ 0 \leq v_j \leq l}} \\ &\quad \times \left\{ \begin{bmatrix} l \\ v_1 \end{bmatrix}_q \dots \begin{bmatrix} l \\ v_D \end{bmatrix}_q q^{\binom{v_1+1}{2} + \dots + \binom{v_D+1}{2}} \gamma_1^{l-v_1} \dots \gamma_D^{l-v_D} \right\}. \end{aligned}$$

By Hilfssatz 4 of [8] it follows that

$$r^{v(l-v)} \begin{bmatrix} l \\ v \end{bmatrix}_q \in \mathbb{Z}, \quad v = 0, 1, \dots, l.$$

If $v_1 + \dots + v_D = i$, then we have

$$q^{-il + \binom{v_1+1}{2} + \dots + \binom{v_D+1}{2} - \binom{i+\lambda}{2}} r^{-v_1(l-v_1) - \dots - v_D(l-v_D)} = s^{-s(i,l)} r^{r(i,l)},$$

where

$$s(i, l) = il + \binom{i+\lambda}{2} - \left(\sum_{j=1}^D v_j^2 + i \right) / 2,$$

$$r(i, l) = \binom{i+\lambda}{2} + \left(\sum_{j=1}^D v_j^2 - i \right) / 2.$$

Since, for all $0 \leq i \leq Dl$, $0 \leq k \leq Dl + \lambda - 1$, $i + k - Dl \geq 0$,

$$s(i, l) - \binom{i+k-Dl}{2} \leq s(i, l) \leq i(l-1/2) + i^2(D-1)/(2D) + \lambda(2i+\lambda)/2 \leq e(l, \lambda),$$

and

$$r(i, l) - \binom{i+k-Dl}{2} \geq \binom{i+\lambda}{2} + \left(\sum_{j=1}^D v_j^2 - i \right) / 2 - \binom{i+\lambda-1}{2} > 0,$$

it follows immediately that all the numbers $\Omega_{l,\lambda} A_{l,\lambda}(1)$ and $\Omega_{l,\lambda} B_{l,\lambda,l}(1)$ are integers.

We obviously have

$$\left| \begin{bmatrix} l \\ v \end{bmatrix}_q \right| \leq c_2^l |q|^{\delta v(l-v)},$$

where $\delta = 0$, if $|q| < 1$, and $\delta = 1$, if $|q| > 1$. In the case $|q| < 1$ we thus obtain, for all $0 \leq i \leq Dl$, $0 \leq k \leq Dl + \lambda - 1$, $i + k - Dl \geq 0$,

$$|\sigma_{i,l} q^{\binom{i+k-Dl}{2} - \binom{i+\lambda}{2}}| \leq c_3^{l+\lambda} |q|^{-1} |q|^{i(l-1/2) - (\min \Sigma + i)/2} \leq c_3^{l+\lambda} |q|^{-e(l,\lambda)},$$

where we have used the notation

$$\Sigma = v_1^2 + \dots + v_D^2, \quad v_1 + \dots + v_D = i.$$

On the other hand, if $|q| > 1$,

$$|\sigma_{i,l} q^{\binom{i+k-Dl}{2} - \binom{i+\lambda}{2}}| \leq c_4^{l+\lambda} |q|^{-il + \binom{i+\lambda-1}{2} - \binom{i+\lambda}{2} \max\{(\Sigma+i)/2 + l - \Sigma\}} \leq c_5^{l+\lambda} |q|^{(i - \min \Sigma)/2} \leq c_5^{l+\lambda}.$$

These considerations imply the inequality

$$\max\{|\Omega_{l,\lambda} A_{l,\lambda}(1)|, |\Omega_{l,\lambda} B_{l,\lambda,t}(1)|\} \leq c_1^{l+\lambda} h^{e(l,\lambda)},$$

which gives Lemma 2.

Next we shall consider the determinant

$$A_{l,\lambda}(z) = \begin{vmatrix} A_{l,\lambda(0)}(z) & B_{l,\lambda(0),1}(z) & \cdots & B_{l,\lambda(0),D}(z) \\ A_{l,\lambda(1)}(z) & B_{l,\lambda(1),1}(z) & \cdots & B_{l,\lambda(1),D}(z) \\ \vdots & \vdots & & \vdots \\ A_{l,\lambda(D)}(z) & B_{l,\lambda(D),1}(z) & \cdots & B_{l,\lambda(D),D}(z) \end{vmatrix}.$$

Our choice of λ is different from the corresponding choice of Stihl [8], and it implies the following useful lemma.

LEMMA 3. If $\lambda(j) = \lambda + j$, $j = 0, 1, \dots, D$, then

$$A_{l,\lambda}(z) = K z^{(D^2+D)(l+1/2)+D\lambda},$$

where $K \neq 0$ is a rational constant.

Proof. Our $A_{l,\lambda}$ is a polynomial of degree

$$\leq Dl + \sum_{j=1}^D (Dl + \lambda + j - 1) = (D^2 + D)l + D\lambda + \binom{D}{2}.$$

The coefficient of $z^{(D^2+D)(l+1/2)+D\lambda}$ is the determinant K ,

$$K = \sigma_{0,l} q^{-\binom{D}{2}} \begin{vmatrix} E_{Dl+\lambda(1)-1,1} & \cdots & E_{Dl+\lambda(1)-1,D} \\ E_{Dl+\lambda(2)-1,1} & \cdots & E_{Dl+\lambda(2)-1,D} \\ \vdots & & \vdots \\ E_{Dl+\lambda(D)-1,1} & \cdots & E_{Dl+\lambda(D)-1,D} \end{vmatrix},$$

where, by Lemma 1,

$$\begin{aligned} E_{Dl+\lambda(j)-1,t} &= \sum_{i=0}^{Dl} \sigma_{i,l} \gamma_t^{i+\lambda(j)-1} q^{\binom{i+\lambda(j)-1}{2} - \binom{i-\lambda(j)}{2}} \\ &= \sum_{i=0}^{Dl} \sigma_{i,l} (\gamma_t/q)^{i+\lambda(j)-1} \\ &= (\gamma_t/q)^{\lambda(j)-1} \sum_{i=0}^{Dl} \sigma_{i,l} (\gamma_t/q)^i \\ &= (\gamma_t/q)^{\lambda(j)-1} \prod_{u=1}^D (\gamma_u - \gamma_t/q, q^{-1})_l, \quad j, t = 1, \dots, D. \end{aligned}$$

Therefore

$$K = (\gamma_1 \cdots \gamma_D)^{l+\lambda} q^{-\binom{\lambda}{2} - D\lambda} \prod_{t=1}^D \prod_{u=1}^D (\gamma_u - \gamma_t/q, q^{-1})_l \\ \times \begin{vmatrix} 1 & \cdots & 1 \\ \gamma_1/q & \cdots & \gamma_D/q \\ \vdots & & \vdots \\ (\gamma_1/q)^{D-1} & \cdots & (\gamma_D/q)^{D-1} \end{vmatrix},$$

and since $\gamma_u \neq \gamma_t q^n$ for all $u \neq t$, $n \in \mathbb{Z}$, we have $K \neq 0$.

On the other hand, since $A_{l,\lambda}(z) f(\gamma_t z) - B_{l,\lambda,t}(z) = R_{l,\lambda,t}(z)$ (see Lemma 1), $\Delta_{l,\lambda}$ can be expressed in the form

$$\Delta_{l,\lambda}(z) = \pm \begin{vmatrix} A_{l,\lambda(0)}(z) & R_{l,\lambda(0),1}(z) & \cdots & R_{l,\lambda(0),D}(z) \\ A_{l,\lambda(1)}(z) & R_{l,\lambda(1),1}(z) & \cdots & R_{l,\lambda(1),D}(z) \\ \vdots & \vdots & & \vdots \\ A_{l,\lambda(D)}(z) & R_{l,\lambda(D),1}(z) & \cdots & R_{l,\lambda(D),D}(z) \end{vmatrix}.$$

Thus $\Delta_{l,\lambda}$ has at $z=0$ a zero of order at least

$$\sum_{j=0}^{D-1} (1 + Dl + \lambda + j) = (D^2 + D)l + D\lambda + \binom{D}{2}.$$

Together with the above considerations this implies Lemma 3.

In the following Section 5 we shall consider more closely the properties of the above used functions $R_{l,\lambda,t}$.

5. THE ESTIMATION OF $R_{l,\lambda,t}$

We also need an upper estimate for the p -value of $\Omega_{l,\lambda} R_{l,\lambda,t}(1)$, $t=1, \dots, D$. This is achieved in the following lemma.

LEMMA 4. *If $p|s$, then the inequalities*

$$|\Omega_{l,\lambda} R_{l,\lambda,t}(1)|_p \leq c_6^{l+\lambda} |s|_p^{e(l,\lambda) + (l^2 + 2l\lambda)/2}, \quad t=1, \dots, D,$$

are valid for all $l \geq c_7$.

Proof. We have, by Lemma 1,

$$\begin{aligned}
 |R_{l,\lambda,t}(1)|_p &\leq \max_{k \geq l+Dl+\lambda} \{|E_{k,t}|_p\} \\
 &\leq \max_{\substack{k \geq l+Dl+\lambda \\ 0 \leq i \leq Dl}} \{|\sigma_{i,t} \gamma_t^{i+k-Dl} q^{\binom{i+k-Dl}{2} - \binom{i+\lambda}{2}}|_p\} \\
 &\leq \max_{0 \leq i \leq Dl} \{|\sigma_{i,t} \gamma_t^{i+l+\lambda} q^{\binom{i+l+\lambda}{2} - \binom{i+\lambda}{2}}|_p\} \\
 &\quad \times \max_{\substack{k \geq l+Dl+\lambda \\ 0 \leq i \leq Dl}} \{|\gamma_t^{k-(l+Dl+\lambda)} q^{\binom{i+k-Dl}{2} - \binom{i+l+\lambda}{2}}|_p\}.
 \end{aligned}$$

Since

$$\binom{i+k-Dl}{2} - \binom{i+l+\lambda}{2} = \sum_{j=i+l+\lambda}^{i+k-Dl-1} j,$$

and $|\gamma_t q^j|_p < 1$, $t = 1, \dots, D$, for all $j \geq c_8$, the last term in the above product is < 1 for all $l \geq c_7$. Therefore

$$\begin{aligned}
 |R_{l,\lambda,t}(1)|_p &\leq c_9^{l+\lambda} \max_{0 \leq i \leq Dl} \{|q|_p^{-il + \min(\Sigma+i)/2} |q|_p^{\binom{i+l+\lambda}{2} - \binom{i+\lambda}{2}}\} \\
 &\leq c_9^{l\lambda} \max_{0 \leq i \leq Dl} \{|q|_p^{(l^2+2l\lambda+\min(\Sigma+i)-l)/2}\} \\
 &\leq c_{10}^{l+\lambda} |s|_p^{(l^2+2l\lambda)/2}
 \end{aligned}$$

for all $l \geq c_7$. By using the definition of $\Omega_{l,\lambda}$ we now immediately obtain Lemma 4.

6. THE PROOF OF THEOREM 1 AND ITS COROLLARY

Let us put

$$L = H_0 + \sum_{t=1}^D H_t f(\gamma_t),$$

where H_t are integers, not all zero, and $\max\{|H_t|\} \leq H$. Then we immediately have

$$\begin{aligned}
 \Omega_{l,\lambda} A_{l,\lambda}(1)m &= \Omega_{l,\lambda}(A_{l,\lambda}(1)H_0 + \sum_{t=1}^D H_t(B_{l,\lambda,t}(1) + R_{l,\lambda,t}(1))) \\
 &= \Omega_{l,\lambda}(A_{l,\lambda}(1)H_0 + \sum_{t=1}^D B_{l,\lambda,t}(1)H_t) \\
 &\quad + \sum_{t=1}^D H_t \Omega_{l,\lambda} R_{l,\lambda,t}(1).
 \end{aligned} \tag{4}$$

Let us choose for λ the value $[\beta l]$, where $\beta > 0$ is a constant that will be fixed later. Then, by Lemma 3, there exists a $\lambda(j) = \lambda + j$, $0 \leq j \leq D$, such that the number

$$Y = \Omega_{l, \lambda(j)}(Z_{l, \lambda(j)} H_0 + \sum_{t=1}^D B_{l, \lambda(j), t}(1) H_t)$$

is different from zero. Further, by Lemma 2, $Y \in \mathbb{Z}$. Thus

$$1 \leq |Y| \mid Y|_p.$$

By using again Lemma 2 we obtain an estimate

$$|Y| \leq H c_{11}^{l + \lambda(j)} h^{e(l, \lambda(j))},$$

which implies an inequality

$$\begin{aligned} 1 &\leq H c_{11}^{l + \lambda(j)} h^{e(l, \lambda(j))} |Y|_p \\ &\leq H c_{11}^{l + \lambda(j)} h^{e(l, \lambda(j))} \\ &\quad \times \max \left\{ |\Omega_{l, \lambda(j)} A_{l, \lambda(j)}(1) L|_{p'} \left| \sum_{t=1}^D H_t \Omega_{l, \lambda(j)} R_{l, \lambda(j), t}(1) \right|_p \right\}. \end{aligned} \quad (5)$$

From Lemma 4 it now follows that for all $l \geq c_7$

$$\left| \sum_{t=1}^D H_t \Omega_{l, \lambda(j)} R_{l, \lambda(j), t}(1) \right|_p \leq c_6^{l + \lambda(j)} |s|_p^{e(l, \lambda(j)) + (l^2 + 2l\lambda(j))/2}.$$

Therefore, let us consider the expression

$$X = H(c_6 c_{11})^{l + \lambda(j)} h^{e(l, \lambda(j))} |s|_p^{e(l, \lambda(j)) + (l^2 + 2l\lambda(j))/2}.$$

Since $\lambda = (\beta l]$, we have

$$X \leq H c_{12}^l (h^{1 - \Gamma(\beta)} |s|_p)^{\Gamma_1(\beta) l^2 / 2},$$

where $\Gamma_1(\beta) = \beta^2 + 2(D+1)\beta + D^2 + D + 1$, $\Gamma(\beta) = (1 + 2\beta)/\Gamma_1(\beta)$. The choice $\beta = \bar{\beta} = (-1 + (1 + 4D^2)^{1/2})/2$ gives a maximal value

$$\Gamma = (2D + 1 - (1 + 4D^2)^{1/2})/(2D)$$

for $\Gamma(\beta)$. Let us denote

$$R = (h^{1 - \Gamma} |s|_p)^{\Gamma_1 / 2}, \quad \Gamma_1 = \Gamma_1(\bar{\beta}).$$

Then $R < 1$, since $h^{1 - \Gamma} |s|_p < 1$ by our assumption (2). With a given $\varepsilon > 0$ we now choose

$$l = [((1 + \varepsilon)(\log H)/\log R^{-1})^{1/2}] + 1.$$

Then $X < 1$ for all $H \geq c_{13}$. Here we suppose c_{13} is large enough to imply $l \geq c_7$. By (5) it therefore follows that

$$1 \leq H c_{11}^{l+\lambda(j)} h^{e(l, \lambda(j))} |\Omega_{l, \lambda(j)} A_{l, \lambda(j)}(1) L|_p. \quad (6)$$

Lemma 2 says that $\Omega_{l, \lambda(j)} A_{l, \lambda(j)}(1) \in \mathbb{Z}$. Thus (2) and (6) give the estimates

$$\begin{aligned} 1 &\leq |L|_p H c_{14}^l h^{\Gamma_1(1-\Gamma)l^2/2} \\ &\leq |L|_p H c_{15}^{(\log H)^{1/2}} H^{(1+\varepsilon)\Gamma_1(1-\Gamma)(\log h)/(-2\log R)} \\ &\leq |L|_p c_{15}^{(\log H)^{1/2}} H^{1-(1+\varepsilon)(1-\Gamma)(\log h)/((1-\Gamma)\log h + \log |s|_p)} \\ &< |L|_p H^{1+(1+2\varepsilon)(1-\Gamma)/(\Gamma-\gamma)}, \end{aligned}$$

for all $H \geq c_{16}$. Here $(1-\Gamma)/(\Gamma-\gamma) = \theta(\gamma, D) - 1 = \theta - 1$. Thus we have

$$|L|_p > H^{-\theta - 2\varepsilon(\theta - 1)}$$

for all $H \geq c_{16}$. This proves Theorem 1.

To prove the corollary it is enough to note that by the choice $q = p^{2a}$ we have

$$g(z) = p^c f(p^{b-a} z).$$

7. THE PROOF OF THEOREM 2

Suppose that

$$L_p = H_0 + \sum_{t=1}^D H_t f_p(\gamma_t) = 0$$

for all $p|s$. As in the beginning of the above proof of Theorem 1 we find again a rational integer

$$Y = \Omega_{l, \lambda(j)}(A_{l, \lambda(j)}(1) H_0 + \sum_{t=1}^D B_{l, \lambda(j), t}(1) H_t) \neq 0.$$

Thus

$$1 = |Y| \prod_p |Y|_p \leq |Y| \prod_{p|s} |Y|_p,$$

where, as before

$$|Y| \leq H c_{11}^{l+\lambda(j)} h^{e(l, \lambda(j))}.$$

We supposed above that $L_p = 0$ for all $p|s$. By (4) it thus follows that

$$|Y|_p = \left| \sum_{t=1}^D H_t \Omega_{l, \lambda(j)} R_{l, \lambda(j), t}(1) \right|_p$$

for all $p|s$. Therefore Lemma 4 implies an inequality

$$|Y|_p \leq c_{17}^{l+\lambda(j)} |s|_p^{e(l, \lambda(j)) + (l^2 + 2l\lambda(j))/2}$$

for all $p|s$, $l \geq c_7$. By choosing λ similar to as in the above proof we now obtain

$$1 \leq H c_{18}^l \left(h^{1-\Gamma} \prod_{p|s} |s|_p \right)^{\Gamma_1 l^2/2}. \quad (7)$$

By the assumption of Theorem 2 we have $(1-\Gamma) \log h < \log |s|$, which implies an inequality $h^{1-\Gamma}/|s| < 1$. Thus (7) leads to a contradiction for all sufficiently large l . This proves Theorem 2.

REFERENCES

1. GEORGE E. ANDREWS, The theory of partitions, in "Encyclopedia of Mathematics and its Applications," Vol. 2, Addison-Wesley, Reading, MA, 1976.
2. J.-P. BEZIVIN, Indépendance linéaire des valeurs des solutions transcendentes de certaines équations fonctionnelles, *Manuscripta Math.* **61** (1988), 103–129.
3. P. BUNDSCHUH AND I. SHIOKAWA, A measure for the linear independence of certain numbers, *Results in Math.* **7** (1984), 130–144.
4. ORHAN S. İÇEN, Über die Funktionswerte der p -adischen Reihe $\sum_{n=0}^{\infty} p^{a \binom{n}{2}} z^n$, *J. Reine Angew. Math.* **202** (1961), 100–106.
5. W. MAIER, Potenzreihen irrationalen Grenzwertes, *J. Reine Angew. Math.* **156** (1927), 93–148.
6. O. PERRON, "Irrationalzahlen," Chelsea, New York, 1951.
7. TH. SKOLEM, Some theorems on irrationality and linear independence, in "Den 11te Skandinaviske Matematikerkongress Trondheim 1949 (1949)," pp. 77–98.
8. THOMAS STIHL, Arithmetische Eigenschaften spezieller Heinescher Reihen, *Math. Ann.* **268** (1984), 21–41.
9. TH. STIHL UND R. WALLISSER, Zur Irrationalität und linearen Unabhängigkeit der Werte der Lösungen einer Funktionalgleichung von Poincaré, *J. Reine Angew. Math.* **341** (1983), 98–110.
10. L. TSCHAKALOFF, Arithmetische Eigenschaften der unendlichen Reihe $\sum_{n=0}^{\infty} a^{-n(n-1)/2} x^n$, *Math. Ann.* **80** (1921), 62–74.
11. K. VÄÄNÄNEN UND R. WALLISSER, Zu einem Satz von Skolem über die lineare Unabhängigkeit von Werten gewisser Thetareihen, *Manuscripta Math.* **65** (1989), 199–212.